



Janice K. Brewer
Governor

Joey Ridenour
Executive Director

Arizona State Board of Nursing

4747 N. 7th Street, Suite 200
Phoenix, AZ 85014-3653
Phone (602) 771-7800 Fax (602) 771-7888
E-Mail: arizona@azbn.gov
Home Page: <http://www.azbn.gov>

An advisory opinion adopted by AZBN is an interpretation of what the law requires. While an advisory opinion is not law, it is more than a recommendation. In other words, an advisory opinion is an official opinion of AZBN regarding the practice of nursing as it relates to the functions of nursing. Facility policies may restrict practice further in their setting and/or require additional expectations related to competency, validation, training, and supervision to assure the safety of their patient population and or decrease risk.

OPINION: Palliative Sedation at End of Life
DATE APPROVED: 1/07
REVISED DATE: 03/10
ORIGINATING COMMITTEE:
SCOPE OF PRACTICE COMMITTEE

Within the Scope of Practice of RN LPN

ADVISORY OPINION PALLIATIVE SEDATION AT END OF LIFE

It is within the scope of practice of a registered nurse to administer medications that provide palliative sedation at the end of life. This may include medications classified as anesthetic agents, but not to be administered to provide an anesthetic as in A.R.S. 32-1661.

For the purpose of this advisory opinion, palliative sedation is defined as follows: the monitored use of medications intended to provide relief of refractory symptoms but not to intentionally hasten death.

A refractory symptom is one that cannot be adequately controlled in a tolerable time frame despite aggressive use of usual therapies, and seems unlikely to be adequately controlled by further invasive or noninvasive therapies without excessive or intolerable acute or chronic side effects/complications.

I. GENERAL REQUIREMENTS

- A. Written Policy and Procedure is maintained by the employer.
 - 1. Administration of medications must be on the order of a person licensed in this state to prescribe such medications.
 - 2. The patient must have sufficient level of nursing care to maintain sedation.
 - 3. The agency has identified medications allowed for palliative sedation, preferably by an interdisciplinary committee including nurses.
 - 4. Pre-sedation symptom assessment is performed by the RN.
 - 5. Post-sedation symptom assessment and ongoing assessments performed by the RN
- B. Only an RN with all of the following criteria are permitted to administer medications for palliative sedation:
 - 1. Current certification in Basic Life Support (BLS).
 - 2. Must possess sufficient knowledge about the issues surrounding the use of palliative sedation to inform patients, families, and other health care providers in making decisions about its use.

3. Documented supervised clinical practice and competency prior to unsupervised administration of palliative sedation medication.

II. RATIONALE

Patients at the end of life may experience physical, psychological, spiritual and existential distress which in most cases can be relieved by optimal end of life care. However, some patients at times may suffer from intractable symptoms. For those patients who are imminently dying and whose suffering is unrelenting and unendurable, palliative sedation may offer relief. The intent of palliative sedation is to relieve suffering in dying patients but not to deliberately hasten death. This distinction separates it from euthanasia and/or assisted suicide where the intent is solely to end life.

III. REFERENCES

- Beel, A., McClement, S.E., Harlos, M. (2002). Palliative sedation therapy: a review of definitions and usage. *International Journal of Palliative Nurse*, 8 (4): 190-9.
- Bruce, S. Hendrix, C. & Gentry, J. (2006). Palliative sedation in end-of-life care. *Journal of Hospice and Palliative nursing*, 8(6), 320-327.
- Claessens, P., Genbrugge, E., Vannuffelen, R., Broeckaert, b., Schotsmans, P. & Menten, J. (2007). Palliative sedation and nursing. *Journal of Hospice and Palliative Nursing*, 9(2), 100-106.
- De Graeff, A. & Mervyn, D. (2007). Palliative sedation therapy in the last weeks of life: A literature review and recommendations for standards. *Journal of Palliative Medicine*, 10(1), 67-85.
- Elsayem, A., Curry, E., Boohene, J., Munsell, M., Calderon, B., Hung, F. & Bruera, E. (2009). Use of palliative sedation for intractable symptoms in the palliative care unit of a comprehensive cancer center. *Support Care in Cancer*, 17(1), 53-59.
- Hospice and Palliative Nurses Association Position Statement, Palliative Sedation at End of Life, www.hpna.org, retrieved May 25, 2006
- Krakauer, E.L., Penson, R.T., Truog, R.D., King, L.A., Chabner, B.A., & Lynch, T.J, Jr. (2000). Sedation for intractable distress of a dying patient: acute palliative care and the principle of double effect. *The Oncologist* 5; 53-62.
- Mercadante, S., DeConno, F., & Ripamonti, C. (1995). Propofol in terminal care. *Journal of Pain and Symptom Management*, 10 (8), 639-642.
- Moyle, J. (1995). The use of propofol in palliative medicine. *Journal of Pain and Symptom Management*, 10 (8), 643.
- Quill, T.E., & Byock, I.R. (2000). Responding to intractable terminal suffering: The role of terminal sedation and voluntary refusal of food and fluids. *Annals of Internal Medicine*, 132 (5): 408-413.

Vermont State Board of Nursing, May, 2007. The Role of the Nurse in the Administration of Propofol (Diprivan) Position Statement.