



Arizona State Board of Nursing

4747 North 7th Street, Suite 200
Phoenix, AZ 85014-3655
Phone (602) 771-7800 Fax (602) 771-7888
E-Mail: arizona@azbn.gov
Home Page: www.azbn.gov

An advisory opinion adopted by AZBN is an interpretation of what the law requires. While an advisory opinion is not law, it is more than a recommendation. In other words, an advisory opinion is an official opinion of AZBN regarding the practice of nursing as it relates to the functions of nursing. Facility policies may restrict practice further in their setting and/or require additional expectations related to competency, validation, training, and supervision to assure the safety of their patient population and or decrease risk.

Within the Scope of Practice of ___ RN ___LPN

**OPINION: INJECTION OF
LIDOCAINE**

APPROVED: 11/89

REVISED DATE: 11/02; 11/02; 1/08, 3/11

ORIGINATING COMMITTEE:

SCOPE OF PRACTICE COMMITTEE

ADVISORY OPINION TUMESCENT LIDOCAINE INJECTION: ROLE OF THE RN

It is **NOT** within the scope of practice of the Registered Nurse (RN) to administer tumescent Lidocaine.

I. RATIONALE

Tumescent Lidocaine Injection is a high risk anesthesia procedure involving large quantities of drug combinations that require the knowledge and skills of a licensed independent provider. It is beyond the knowledge and skills of the RN creating a significant risk to patient safety including possible loss of life.

II. REFERENCES

Clayman, M. A., & Seagle, B. M. (2006, Sept.). Office Surgery safety: The myths and truths behind the Florida Moratoria—six years of Florida data. *Plastic & Reconstructive Surgery*, 118 (3), 777-785.

Commons, G. W., Halperin, B., & Chang, C. C. (2001, Nov.). Large-volume liposuction: A review of 631 consecutive cases over 12 years. *Plastic & Reconstructive Surgery*, 108 (6), 1753-1763.

De Jong, R. H. (2005, January). "Tumescent anesthesia" for office based liposuction. *Anesthesia & Analgesia*, 100 (1), 299-300.

Habbema, L. (2009, July). Safety of liposuction using exclusively tumescent local anesthesia in 3, 240 consecutive cases. *Dermatologic Surgery*, 35 (11), 1728-1735.

Lehnhardt, M., Homann, H. H., Daigeler, A., Hauser, J., Palka, P., & Steinau, H. U. (2008, June). Major and lethal complications of liposuction? A review of 72 cases in Germany between 1998 and 2002. *Plastic & Reconstructive Surgery*, 121 (6), 396e-403e.

Nordstrom, H., & Stange, K. (2005, Oct.). Plasma lidocaine levels and risks after liposuction with tumescent anesthesia. *Acta Anaesthesiologica Scandinavica*, 49 (10), 1487-1490.

Salim-Tabbousch, Z. (2004, April). Tumescent anesthesia: A concern for anesthesiologist. *Anesthesia & Analgesia*, 98 (4), 1190.